

2018-03-20

# Specific protein supplementation using soya, casein or whey differentially affects regional gut growth and luminal growth factor bioactivity in rats; implications for the treatment of gut injury and stimulating repair

Marchbank, Tania

<http://hdl.handle.net/10026.1/10365>

---

10.1039/c7f01251a

Food and Function

Royal Society of Chemistry

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

**Specific protein supplementation using soya, casein or whey differentially affects regional gut growth and luminal growth factor bioactivity in rats; implications for the treatment of gut injury and stimulating repair**

Tania Marchbank<sup>1,2</sup>, Nikki Mandir<sup>3</sup>, Denis Calnan<sup>4</sup>, Robert A. Goodlad<sup>3</sup>, Theo Podas<sup>4</sup>, Raymond J. Playford<sup>1</sup>

1. Plymouth University Peninsula Schools of Medicine and Dentistry, Plymouth, UK.  
2. Centre of Immunobiology, Blizard Institute, Barts and The London School of Medicine, Queen Mary, University of London, UK. 3. Histopathology, CRUK, Lincoln's Inn Fields Laboratory, 44 Lincoln's Inn Fields, London, UK. 4. University Division of Gastroenterology, Leicester General Hospital, Gwendolen Road, Leicester, UK.

**Correspondence** Prof. RJ Playford, Peninsula Medical School, Plymouth University, The John Bull Building, Tamar Science Park, Research Way, Plymouth PL6 8BU

**Tel:** +441752582002

**Email:** raymond.playford@plymouth.ac.uk

**Running title:** Specific dietary protein supplementation and regional gut growth

**Abbreviations:** ANOVA; analysis of variance, C; casein, ED; elemental diet, EGF; epidermal growth factor, GLP-2; glucagon-like peptide-2, SCFAs; short chain fatty acids, SI; small intestine, SP; soya protein, W; whey protein,

25

26 **ABSTRACT**

27 Modulation of regional growth within specific segments of the bowel may have  
28 clinical value for several gastrointestinal conditions. We therefore examined the  
29 effects of different dietary protein sources on regional gut growth and luminal growth  
30 factor bioactivity as potential therapies. Rats were fed for 14 days on isonitrogenous  
31 and isocaloric diets comprising elemental diet (ED) alone (which is known to cause  
32 gut atrophy), ED supplemented with casein or whey or a soya protein-rich feed.  
33 Effects on regional gut growth and intraluminal growth factor activity were then  
34 determined. Despite calorie intake being similar in all groups, soya rich feed caused  
35 20% extra total body weight gain. Stomach weight was highest on soya and casein  
36 diets. Soya enhanced diet caused greatest increase in small intestinal weight and  
37 preserved luminal growth factor activity at levels sufficient to increase proliferation *in*  
38 *vitro*. Regional small intestinal proliferation was highest in proximal segment in ED  
39 fed animals whereas distal small intestine proliferation was greater in soya fed  
40 animals. Colonic weight and proliferation throughout the colon was higher in animals  
41 receiving soya or whey supplemented feeds. We conclude that specific protein  
42 supplementation with either soya, casein or whey may be beneficial to rest or increase  
43 growth in different regions of the bowel through mechanisms that include  
44 differentially affecting luminal growth factor bioactivity. These results have  
45 implications for targeting specific regions of the bowel for conditions such as Crohn's  
46 disease and chemotherapy.

47

47

48 **INTRODUCTION**

49 Many intrinsic inflammatory and ingested injurious agents cause regional damage to  
50 the gut. These include Crohn's disease (mainly affecting distal small intestine),  
51 ulcerative colitis (affecting colon) and non-steroidal inflammatory drugs (mainly  
52 affecting stomach and small intestine). Therapeutic targeting of drugs may occur  
53 through site specific release formulations or topical therapy (such as colonic enemas).  
54 As an alternative to pharmacological approaches, there is currently interest in using  
55 nutritional manipulation to both reduce the risk of relapse and for treatment when an  
56 injury is present. For example, elemental diets (ED), consisting of amino acids,  
57 mono/disaccharides and short/medium chain fatty acids, are of proven value in the  
58 treatment of inflammatory bowel disease, especially in children.<sup>1,2</sup> The mechanism(s)  
59 of action of EDs are unclear but might include reducing systemic antigenic load and  
60 "resting" the bowel. More recently, polymeric diets have been shown to be equally  
61 efficacious to ED<sup>3</sup>, establishing the principal that selective additions to feeds may  
62 enhance efficacy without losing the beneficial effects of ED. Prolonged use of ED  
63 causes intestinal atrophy, particularly of the distal bowel, potentially increasing the  
64 risk of bacterial translocation and bacteraemia.<sup>4</sup> Dietary formulations which reduce  
65 gut luminal antigenic load whilst minimizing the associated gut atrophy could,  
66 therefore, have therapeutic advantages. Similarly, because the gut is second only to  
67 the bone marrow in cell turnover, damage to the intestine is a common issue in  
68 patients undergoing chemotherapy (chemotherapy induced mucositis) and diets which  
69 reduce turnover while chemotherapy is administered or enhancing growth after  
70 treatment may also be useful.

The physiological control of gut growth is complex and poorly understood. Possible controlling mechanisms include alteration in circulating trophic hormones, local nutrition and luminal growth factors. We have previously suggested that ingestion of specific food proteins could influence gut growth by reducing the destruction of luminal growth factors, such as epidermal growth factor (EGF), through acting as competitive substrates for pancreatic proteases.<sup>5</sup> In this model, the efficacy of the various proteins to stimulate gut growth would be dependent on their ability to act as substrate for pancreatic proteases, relative to the affinity of the luminal growth factors for the same enzymes. Support for this idea comes from an *in vitro* study that showed soya bean trypsin inhibitor or casein were better substrates than whey protein (“lactalbumin”) for human pancreatic proteases. Furthermore, the addition of soya protein or casein to human duodenal juice reduced its ability to digest EGF, whereas addition of whey protein or elemental diets did not.<sup>5</sup> Soya protein or casein supplementation of feeds such as elemental diets might, therefore, be a clinically useful method of decreasing gut hypoplasia, while having a relatively minor effect on “antigenic load”.

We, therefore, examined the effect of isocaloric and isonitrogenous diets comprising standard elemental diet (E028, Nutricia, Liverpool, UK), E028 supplemented with either casein (ED+C) or whey protein (ED+W), and a commercially available soya-protein rich diet (SP, containing human food grade soya bean concentrate) on regional gut growth in rats and also assessed total “growth factor” bioactivity within the lumen of the small intestine.

## **MATERIALS AND METHODS**

96

97 Diets

98 Three of the four test diets comprised standard elemental E028 ('ED', Nutricia  
99 Liverpool, UK) or variants of E028 containing whey protein ('ED+W', 12.5g/100g,  
100 the two major components of whey being  $\alpha$ -lactoglobulin and  $\beta$ -lactoglobulin.)<sup>6</sup>  
101 or casein ('ED+C', 10.5g/100g). Both protein supplemented elemental diets were  
102 specifically produced for the studies by Nutricia Liverpool. The fourth group  
103 consisted of a commercially available soya protein rich diet ('SP', produced by SDS,  
104 Essex, UK). All four diets were similar in terms of digestible protein equivalence and  
105 amino-acid content but the SP diet had a higher carbohydrate content, lower added fat  
106 and contained some crude fibre (5.3%). Details of the various diets are shown in  
107 Supplemental Table 1.

108

109 Protease inhibitory activity of feeds

110 To determine trypsin inhibitory activity within the feeds, a standard BAPNA substrate  
111 assay<sup>7</sup> was undertaken with concentrations of trypsin (0 - 50  $\mu$ g) added to generate a  
112 standard curve. Additional tubes, each containing 20  $\mu$ g trypsin had varying  
113 concentrations of the feeds added and any effect on reduction of tryptic activity (due  
114 to competitive inhibition) determined. In addition, to reproduce the situation found *in*  
115 *vivo*, where the feeds would be exposed to gastric juice prior to entering the small  
116 intestine, additional samples were pre-incubated with pepsin (1mg/ml) for 1 h. at pH 2,  
117 37°C before neutralization and subsequent BAPNA analyses.

118

119 **Animal Experiments**

120 All experiments were performed in compliance with relevant United Kingdom laws

and guidelines. All animal experiments were approved by the Local Institutional Animals Ethics Committees (Imperial College and Cancer Research UK) and performed under institutional guidelines. Studies were covered by the appropriate licences under the United Kingdom Home Office Animals Procedures Acts, 1986 and performed under the UK home office Animal Research: Reporting of In Vivo Experiments) guidelines (ARRIVE).

Pilot Study: We initially examined the caloric intake of rats (5 per group) fed on the various diets if allowed access *ad libitum*. Dietary intake was similar with all of the ED groups but was 10% greater in SP diet fed animals (data not show). Dietary restriction of SP fed animals was, therefore, used for the main study to ensure equal caloric intake between the various groups.

Main study: Forty adult male, Sprague-Dawley rats (Charles River, UK, 180-225g) were coded, weighed and randomized into the four diet groups (n=10 per group) and fed exclusively on these diets for 14 days. Animals remained in their wire bottomed cages in groups of 5 throughout the study. Food was given in 2-ring concentric aluminium containers designed to minimize food spillage. The food consumption of each cage with determined each day by back weighing, including the small amount of spilled food beneath the cages.

#### Autopsy procedure and collection of samples.

At the end of the 14 day study, rats were given vincristine sulphate by intraperitoneal injection (1mg/kg, Eli Lilly, Basingstoke, UK) and killed 2 h later. Analyses of tissue was performed using the methods published by our group previously.<sup>8</sup> The weight and

unstretched length of the various portions of the gastrointestinal tract were measured. To assess the effect of these various diets on luminal growth factor activity, 1 ml of iced, normal saline was then lavaged through the small intestine, collected into Eppendorf tubes, snap frozen using frozen CO<sub>2</sub> (dry ice) and stored at -70 °C until assay for growth factor activity.

The stomach and samples of the small intestine and colon were then fixed in Carnoy's fluid, and stored in 70% (volume/volume) ethanol. For the intestinal tissues, the position of the various samples was defined by expressing its harvest site as a percentage of that organs total length at 1.5, 50 and 90% small intestinal distance and can be considered as equivalent to duodenum, jejunum and ileum respectively. Similarly, 10, 50 and 90% colon can be considered as proximal, mid and distal colon.

#### Analysis of tissue samples.

Assessment of metaphase accumulation was performed on micro-dissected tissue using our well validated, previously published methods<sup>8</sup> by a person blinded to the experiment. Briefly, tissues were sequentially rehydrated, hydrolyzed and stained with Schiff's reagent (the Feulgen reaction) and transferred to 45% (volume/volume) acetic acid. The crypts were then teased apart under a dissecting microscope, transferred to a glass microscope slide, flattened gently beneath a coverslip and examined using a compound microscope. The number of blocked mitoses per crypt was counted and the mean for 20 crypts per animal per site determined and used in the subsequent one way analysis of variance (ANOVA).

#### Analysis of luminal growth factor activity.



Analyses of luminal growth factor activity were determined using thymidine incorporation into primary rat hepatocytes as a bioassay and an EGF-standard curve as described previously by us.<sup>9,10</sup> Briefly, to prepare primary rat hepatocytes for the *in vitro* assay of luminal growth factor activity, male Sprague-Dawley rats were anaesthetized using Hypnorm (Janssen Pharmaceutica-Crown Chemicals Ltd, Lamberhurst, Kent, UK) and hepatocytes isolated by *in situ* collagenase perfusion. The basic protocol consists of a two-step perfusion of the liver *in situ*, via the portal vein, first with calcium-free buffer followed by a calcium supplemented buffer containing collagenases. The digested liver was removed, the cells dispersed, filtered, collected by centrifugation and resuspended in a plating medium. For all studies, hepatocytes were grown in Williams E medium without L-glutamine (Gibco BRL, Paisley, UK) containing 5% fetal calf serum. Cell viability, determined by the ability to exclude 0.2% trypan blue, was greater than 80% in all experiments.

Assay of growth factor activity: Various volumes (50 to 200  $\mu$ l) of luminal perfusate was added to the wells. To assess the percentage of cells entering DNA synthesis, [<sup>3</sup>H] thymidine (2 mCi/well, 10 ml, Amersham International, Bucks, UK) was included in the cultures eight hours after the addition of test samples. The amount of [<sup>3</sup>H] thymidine incorporated was assessed biochemically, 18 hours after the addition of [<sup>3</sup>H] thymidine. Cells were washed for 15 seconds with water using a Dynatech multimash automatic self-harvester and solubilized by incubation at 37°C for 1 h in 200  $\mu$ l of 1 M KOH. 50  $\mu$ l of the cell extract was counted in a  $\beta$ -counter in 1 ml of scintillant (Optiphase safe, LKB-Pharmacia Wallac).

The increase in thymidine uptake was then expressed in terms of EGF-like bioactivity by determining the equivalent amount of EGF that was needed to be added to the hepatocytes, to stimulate a similar increase in thymidine uptake. Measurements of EGF-like bioactivity in intestinal juice samples were performed in quadruplicate in 4 separate wells.

## **Statistical Analysis.**

Data were analysed using one-way ANOVA with the different diets as the factor. Where a significant effect of diet was seen, individual comparisons were performed comparing against ED, using t-tests based on the residual and degrees of freedom obtained from the ANOVA, a method equivalent to repeated measures analysis. All data is expressed as mean +/- SEM unless stated.

## **RESULTS**

### Protease inhibitory activity of feeds

*In vitro* assay of feeds showed higher trypsin inhibitory activity (136 mg trypsin inhibition per 100g feed) in SP feed, moderate amount in ED+C (42 mg trypsin inhibition per 100g feed) but much lower amounts in ED (13 mg/100g) or ED+W (15 mg/100g). Samples which had been pre-incubated with pepsin at pH 2 to reproduce prior exposure to gastric contents gave similar results (SP 120 mg trypsin inhibition per 100g feed, ED+C 38 mg/100g, ED 13 mg/100g and ED+W 12 mg/100g).

### **Animal study**

Dietary intake: Daily caloric consumption per cage for each of the groups were as follows ED; 696, 666 - 726 cal. (mean, range), ED+W; 718, 682 - 759 cal., ED+C;

714, 695 - 742 cal., SP; 705 cal (no range as all food eaten each day). This gave an average daily calorie consumption of 139 per animal in the ED group, 144 in ED+W group, 144 in ED+C group and 141 in SP fed animals. The caloric consumption of all groups were, therefore, within 3% of each other ( $p = 0.085$ ) although because of the different constituents of the SP diet, this resulted in SP fed animals receiving approximately 18% greater protein equivalent intake.

Total body weight gain: All animals increased weight over the two-week test period. with ED group showing an average of  $54.5 \pm 4.4$  g increase. Compared to the ED diet, weight gain in ED+C and ED+W ( $48.5 \pm 3.2$  g and  $58.5 \pm 2.1$  g, respectively) was not significantly different but was 20% greater in SP group ( $65.5 \pm 3.7$  g,  $p < 0.01$ ).

Stomach weight: Absolute stomach weights were similar in ED and ED+W groups but about 10% greater in the ED+C and SP groups ( $p < 0.05$  and  $0.01$  vs. ED, respectively, Figure 1A).

Small intestine (SI) weight, proliferation and intraluminal growth factor assay: Animals fed on the elemental diets all had similar SI weight whereas SP group small intestinal weight was 17% heavier ( $p < 0.01$ , Figure 1B). Correspondingly, luminal growth factor activity equivalent to  $3.9 \pm 0.4$  ng of EGF was present in SP group but unrecordable in all other groups.

Small intestinal regional proliferation varied dependent on site with the different diets. Proximal small intestinal proliferation was greatest in the ED, with the other three groups being significantly lower (Figure 2A). In contrast, SP group had the highest

proliferation rate in the mid small intestine (significantly greater than ED+W and ED+C, Figure 3B) and this difference was particularly marked in the distal small intestine with SP having proliferation rate of 150% of the other diets (Fig 2C).

Colon weight and proliferation: Compared to ED fed animals, total colonic weight was 15% higher in ED+W animals and 50% higher in SP animals (both  $p < 0.01$ , Figure 1C). Regional proliferation data was similar throughout the colon with ED+W and SP animals having 50% (proximal colon), 80% (mid colon) and 120% (distal colon) higher proliferation compared to ED or ED+C groups ( $p < 0.01$ , Figure 3).

## DISCUSSION

We have shown different dietary protein sources cause differential regional effects on gut growth and proliferation and that diets containing soya protein result in increased small intestinal luminal growth factor activity.

Care was taken to ensure caloric intake of the groups were comparable to remove a potential confounding factor. We used fourteen days dietary intervention to reflect what would likely be used in a clinical setting in situations such as pre- or post- chemotherapy intervention. We used a rat hepatocyte system to assay intraluminal growth factor activity as this is a robust, reproducible method to determine relative bioactivity of EGF-R type ligands in the gut lumen and has the advantage over most cell lines of maintaining viability in the presence of bile and other luminal contents.<sup>9</sup> 10 The *in vitro* trypsin (inhibition) assay is a simple robust system to examine differential effects of food substances and builds on our earlier *in vitro* work<sup>5</sup> showing that soya-bean trypsin inhibitor causes marked trypsin inhibition, casein causes

moderate trypsin inhibition (probably acting as a competitive substrate), whereas ED or whey have very limited inhibitory activity.

The trophic effects of the various diets differed dependent on which region of the gut was analyzed. Compared to ED diet, stomach weight was better maintained in ED+C and SP rats. The effect of ED+C may be due to casein being precipitated in stomach acid, delaying gastric emptying, whereas rapid transit into the small intestine is seen in ED or ED+W animals.<sup>11</sup>

The most notable finding of the ED diet was the much higher proliferation rate seen in the proximal small intestine compared to the other diets. This is probably as a result of rapid transit into the proximal intestine, where the ED constituents are rapidly adsorbed. This finding could be explained by the luminal workload hypothesis that suggests that regions actively involved in digestion/absorption have increased proliferation. This would also explain the low mitotic level seen in more distal regions of the small bowel as the majority of the feed would have been adsorbed more proximally.

The most notable effects of ED+W were the increased weight and proliferation of the colon. Several potential mechanisms might have relevance in explaining these findings. Changes in circulating trophic factors such as glucagon-like peptide-2 (GLP-2) may have relevance although it is noteworthy that the trophic effect was restricted to the colon and not the small intestine. GLP-2 is produced by the ileum and colon, is trophic to the small intestine and colon and is rapidly degraded by tissue DPP-IV (dipeptidyl peptidase 4). Liu and co-workers showed oral whey

supplementation decreased colonic tissue DPP-IV in rats receiving continuous intravenous GLP-2, possibly increasing local colonic tissue GLP-2 levels.<sup>12</sup> Therefore, even if circulating levels of GLP-2 remain stable, local changes in DPP-IV as a result of dietary manipulation may alter local concentrations. A further mechanism that would explain our results is our finding that whey proteins are poor substrates for luminal proteases when compared against casein or soya. The amount of nutrient reaching the terminal bowel is, therefore, likely to be higher in the ED+W compared to the other groups. Support for this idea comes from recent studies showing whey supplementation increased colonic short chain fatty acids (SCFAs) concentrations, a known substrate for colonocytes.<sup>13</sup> Further studies to determine the importance of colonic bacteria in maintaining the growth of the colon in these animals could include measurement of colonic SCFAs or use of 'germ free' animals.

The most notable effects of the SP diet in the current studies were the increased total body weight gain, the increased small and large bowel growth and the finding that bioactive growth factor activity was present in small intestinal washings. The additional increase in body weight in the SP-fed group was not due to differences in total calorie intake as these were well matched across the groups although it should be noted that the SP group received a slightly higher percentage of their calories from protein. We have previously suggested that control of intestinal growth might be mediated by changes in intraluminal growth factor concentrations, with the trophic effect dependent on the relative affinity of the luminal proteases for the ingested food and the luminal growth factors.<sup>5</sup> This idea was supported by our findings that pancreatic diversion away from the gut lumen increases luminal growth factor concentrations and caused associated regional hypertrophy.<sup>5</sup> The current studies

extend these findings showing that ingestion of soy based proteins (which includes soya bean trypsin inhibitor) are effective competitive substrates for luminal proteases *in vivo*, allowing luminal growth factors such as EGF to survive better than when animals are fed whey or casein. These findings may have relevance to the trophic effect on the small and large intestine although it must be noted that the SP diet contained some fibre which may have relevance for colonic growth through acting as a fuel source for colonic bacteria, increasing colonic SCFAs levels in a similar way to that discussed for the whey fed animals. Further studies examining ED supplemented with pure soya protein which has had the fibre further reduced, combined with measurement of colonic SCFAs and possibly the use of germ-free animals could help address this question.

There is currently much interest in the use of dietary proteins for the prevention and treatment of ill health.<sup>14</sup> Our findings that dietary manipulation can cause regional changes in gut growth and also modulates luminal growth factor levels has potential clinical relevance. ED or polymeric diets are of proven benefit for Crohn's disease, a condition that predominantly affects the small intestine. Specific protein supplementation of ED with whey may have advantages in "resting" the small intestine while preserving colonic growth, which among other advantages should reduce the risk of bacterial translocation. Supplementation of ED with soy protein may have the advantage of locally increasing luminal EGF levels, facilitating repair in a similar way to that shown by us in treating ulcerative colitis with topical therapy.<sup>15</sup> Further studies examining the effect of these proteins on intraluminal growth factor concentrations and injury and repair in animal models of inflammatory bowel disease appear warranted but go beyond the scope of this paper. Similarly, chemotherapy

344 induced injury to the gut is a common side effect of high dose chemotherapy and  
345 systemic administration of several different growth factors to promote repair have  
346 been tested or are being assessed.<sup>16</sup> Dietary manipulation to reduce gut cellular  
347 turnover prior to treatment of non-gut tumours using systemic chemotherapy (to  
348 reduce gut side effects) and/or to increase gut proliferation post treatment, therefore,  
349 has potential therapeutic advantage by locally increasing luminal growth factor  
350 concentrations without the risk of systemic administration. Further studies appear  
351 justified.



**Funding:** Funded by institutional funding.

**Conflict of Interest Statement:** All authors declare no conflict of interest.

## REFERENCES

1. J. Lee, R. Allen, S. Ashley, S. Becker, P. Cummins, A. Gbadamosi, O. Gooding, J. Huston, J. Le Couteur, D. O'Sullivan, S. Wilson, M.C. Lomer. Dietetic Association evidence-based guidelines for the dietary management of Crohn's disease in adults. *J Hum Nutr Diet.* 2014;**27**,207-218.

2) A.K. Akobeng, V. Miller, J. Stanton, A.M. Elbadri, A.G. Thomas. Double-blind randomized controlled trial of glutamine-enriched polymeric diet in the treatment of active Crohn's disease. *J Pediatr Gastroenterol Nutr.* 2000;**30**,78-84.

3) S. Verma, S. Brown, B. Kirkwood, M.H. Gjafer. Polymeric versus elemental diet as primary treatment in active Crohn's disease: a randomized, double-blind trial Polymeric vs Elemental Diet in Crohn's Disease. *The American Journal of Gastroenterology* 2000;**95**,735-739

4). E.A. Deitch, D. Xu, M.B. Naruhn, D.C. Deitch, Q. Lu, A.A. Marino. Elemental diet and IV-TPN-induced bacterial translocation is associated with loss of intestinal mucosal barrier function against bacteria. *Ann Surg* 1995;**221**,299-307.

5) R.J. Playford, A.C. Woodman, P. Clark, P. Watanapa, D. Vesey, P. Deprez, R.C. Williamson, J. Calam. Effect of luminal growth factor preservation on intestinal growth. *Lancet.* 1993;**341**,843-8.

6) D.P. Mohanty, S. Mohapatra, S. Misra, P.S. P.S. Sahu, Milk derived bioactive peptides and their impact on human health - A review. *Saudi J Biol Sci.* 2016;**23**,577-

- 375 7) T. Marchbank, A. Mahmood, R.J. Playford. Pancreatic secretory trypsin inhibitor  
376 causes autocrine-mediated migration and invasion in bladder cancer and  
377 phosphorylates the EGF receptor, Akt2 and Akt3, and ERK1 and ERK2.  
378 *Am J Physiol Renal Physiol.* 2013,**305**,F382-9.
- 379 8) P.A. Kitchen, R.A. Goodlad, A.J. FitzGerald, N. Mandir, M.A. Ghatei, S.R. Bloom,  
380 J. Berlanga-Acosta, R.J. Playford, A. Forbes, J.R. Walters. Intestinal growth in  
381 parenterally-fed rats induced by the combined effects of glucagon-like peptide 2 and  
382 epidermal growth factor. *J Parenter Enteral Nutr.* 2005,**29**,248-254.
- 383 9) R.J. Playford, T. Marchbank, D.P. Calnan, J. Calam, P. Royston, J.J. Batten, H.F.  
384 Hansen. Epidermal growth factor is digested to smaller, less active forms in acidic  
385 gastric juice. *Gastroenterology* 1995,**108**,92-101.
- 386 10) D.P. Calnan, A. Fagbemi, J. Berlanga-Acosta, T. Marchbank, T. Sizer, K. Lakhoo,  
387 A.D. Edwards, R.J. Playford. Potency and stability of C terminal truncated human  
388 epidermal growth factor. *Gut.* 2000,**47**,622-7.
- 389 11) M. Dangin, Y. Boirie, C. Garcia-Rodenas, P. Gachon, J. Fauquant, P. Callier, O.  
390 Ballèvre, B. Beaufrère. The digestion rate of protein is an independent regulating  
391 factor of postprandial protein retention. *Am J Physiol Endocrinol Metab*  
392 2001,**280**,E340-8.
- 393 12) X. Liu, S.G. Murali, J.J. Holst, D.M. Ney. Whey protein potentiates the  
394 intestinotrophic action of glucagon-like peptide-2 in parenterally fed rats. *Am J*  
395 *Physiol Regul Integr Comp Physiol.* 2009,**297**,R1554-62.
- 396 13) K. Tomoda, K. Kubo, K. Dairiki, T. Yamaji, Y. Yamamoto, Y. Nishii, A.  
397 Nakamura, M. Yoshikawa, K. Hamada, H. Kimura. Whey peptide-based enteral diet  
398 attenuated elastase-induced emphysema with increase in short chain fatty acids in  
399 mice. *BMC Pulm Med.* 2015,**15**,64.

- 14) A. Pihlanto , P. Mattila , S. Mäkinen , A.M. Pajari. Bioactivities of alternative protein sources and their potential health benefits. *Food Funct.* 2017 Aug 14. [Epub ahead of print]
- 15) A. Sinha, J. Nightingale, K.P. West, J. Berlanga-Acosta, R.J. Playford. Epidermal Growth Factor Enemas with Oral Mesalamine for Mild-to-Moderate Left-Sided Ulcerative Colitis or Proctitis. *N Engl J Med* 2003;**349**,350-357.
- 16) C.J. Xian. Roles of growth factors in chemotherapy-induced intestinal mucosal damage repair. *Curr Pharm Biotechnol* 2003;**4**,260-269.

## FIGURE LEGENDS

### **Figure 1. Effect of diets on percentage change in gastrointestinal weight**

Animals were fed for two weeks on isonitrogenous feeds comprising elemental diet (ED) alone, ED supplemented with milk whey (ED+W), ED supplemented with casein (ED+C) or fed a commercial soya protein rich diet (SP). All groups received similar total calorie intake. Changes in weight of the stomach (**A**), small intestine (**B**) and colon (**C**) after the two-week period were expressed as % change compared to that of animals fed ED diet. \* and \*\* signifies  $p < 0.05$  and  $p < 0.01$ , respectively compared with ED.

### **Figure 2. Effects of diets on small intestinal mitosis.**

All animals on the various diets were injected with vincristine two hours prior to killing to allow assessment of rate of mitosis. Tissue was microdissected and data expressed as number of mitosis per crypt. The three regions analyzed were located according their position as a percentage of total small intestinal length; proximal small intestine (1.5%, duodenum, **A**), mid small intestine (50%, jejunum, **B**) and distal small intestine (90%, ileum, **C**). \* and \*\* signifies  $p < 0.05$  and  $p < 0.01$ , respectively

compared with ED.

**Figure 3. Effects of diets on colon mitosis.**

Colonic tissue from the same animals as Fig 3 was collected and the number of mitosis per crypt assessed. The regions analyzed were located according their position as a percentage of total colonic length; 5% site of colon (proximal colon, **A**), 50% site of colon (mid-colon, **B**), 90% site of colon (distal colon, **C**). \* and \*\* signifies  $p < 0.05$  and  $p < 0.01$ , respectively compared with ED.




Fig 1




Fig 2

434

435

436




Fig 3

437

438

439

440

441 **Supplemental Table 1. Composition of feeds used for study.**

	E028 (ED)	E028 and casein 400g cans (ED + C)	E028 and whey protein 400g cans (ED + W)	Soya protein enhanced diet (SP)
Digestible protein equivalent (g/100g)	12.5	13.8	13.8	12.9
Energy content (kcal/100g)	443	447	447	354
Total amino acid content (g)	15	15	15	14.22
Fat content (g)	17.45	17.45	17.45	2.6
Carbohydrate source	Dried glucose syrup	Dried glucose syrup	Dried glucose syrup	Wheat, barley,
Nitrogen source	Crystalline amino acids	Crystalline amino acids +  casein proteins	Crystalline amino acids + whey proteins	Soya protein, soya protein concentrate, wheat and barley proteins
Fat source	Coconut, safflower and canola oil	Coconut, safflower and canola oil	Coconut, safflower and canola oil	Soya oil
Nitrogen content (%)	2.2	2	2.1	2.35

442

443 1. ED – elemental diet

444 2. ED + C - elemental diet + casein

445 3. ED + W - elemental diet + whey protein

446 4. SP-soya protein enhanced diet

447

448